FORAGE CROPS SELECTION: PROGRESS AND CHALLENGES

Z.SH. SHAMSUTDINOV

W.R. Williams All-Russian Fodder Research Institute, Russian Academy of Agricultural Sciences, korp. 1, ul. Nauchnii Gorodok, Lobnya, Moscow Province, 141055 Russia, e-mail aridland@mtu-net.ru
Received January 30, 2013

Abstract

In the Breeding Center of V.R. Williams All-Russian Fodder Research Institute more than 150 varieties of forage plants have been created for 42 years. Of these, the most widespread 85 varieties cultivated in Russia are not inferior to the best foreign varieties in productivity and exceed them in hardiness, the edaphic resistance to acidity and salinity, and phytocenotic compatibility in mixtures. The *Trifolium pratense* L. varieties VIK 7, tetraploid VIK, Altyn, Topaz, Rannii 2, Trio, Mars, and the *T. repens* L. varieties Yubileynii, VIK 70 are characterized by a precisely expressed ecological individuality, symbiotic activity and different maturing time. The *Medicago* L. new varieties Vega 87, Lada, Nakhodka, Pastbishchnaya 88, Lugovaya 67, Selena, Soleustoichivaya, with the dry matter yield at 12-14 t/ha in the Central non-Chernozem zone, possess various phytocenotic, edaphic, symbiotic properties. More than 40 new varieties of *Bromopsis inermis* Leys., *Phleum pratense* L., *Festuca pratensis* Huds, *Dactylis glomerata* L., *Lolium perenne* L., *Poa pratensis* L., providing yield of 11-12 t/ha and above, are characterized by improved fodder quality and resistance to diseases. In recent years, the varieties of *Vicia sativa* L. and *V. villosa* Roth., high-resistant to environmental and phytocenotic factors, are created for different use. The *V. sativa* L. varieties Lugovskaya 48, Uzunovskaya 91, Vera, and the *V. villosa* Roth. variety Lugovskaya express precocity and a reduced insistence to heat, especially during fruiting. For restoration of lost biodiversity and efficiency of the degraded pastures in arid areas of the Russian South, 18 new varieties of fodder xerohalophytes are created, particularly Barkhan, Dzhangar of *Kochia prostrata* (L.) Schrad., Salang of *Salsola orientalis* S.G. Gmel., Nogana, Alsu of *Camphorosma lessingii* (Litv.), Favorite, Tulkin, Bar of *Eurotia ceratoides* (L.) C.A. Mey., etc. These varieties are widely used for restoration of deserted lands. With reference to perennial bean and cereal grasses, more than 25 breeding technologies are patented in the Russian Federation. Adaptive systems for seed reproduction and the «know-how» for growing perennial and annual grasses are offered. With regard to the achieved results and recent data of fundamental biology a new breeding paradigm is substantiated, based on biogeocenotic principles.

Keywords: breeding, seed, fodder crops, breeding techniques, alfalfa and perennial grasses, arid forage plants.

December 2014 marks the 42nd anniversary of the founding the selection center as part of the W.R. Williams All-Russian Fodder Research Institute. Formed in the Soviet Union the complex for breeding and seed production of fodder crops currently has fairly high scientific potential of 6 specialized and 12 complex breeding centers and more than 20 scientific breeding and seed production units in research institutions. The Breeding Center of the W.R. Williams All-Russian Fodder Research Institute is leading in the development of theory and practice of seed selection.

The main fundamental and applied aspects the researchers of the institute are working on are as follows: i) creating system of climatically and environmentally differentiated and economically specialized fodder crops complementary each other; ii) development and improvement of breeding and selection technologies; iii) original and commercial seed production; iii) education; iii) coordinating national and international programs on fodder crops.

In the W.R. Williams All-Russian Fodder Research Institute there were created more than 150 varieties of clover, alfalfa, annual leguminous fodder crops, perennial cereal grasses, arid fodder plants of new generation, including 85 varieties for past 20 years. The progress in fodder grass production in most

Russia regions, particularly in the humid zone, is most due to clover breeding. The studies of biological, ecological and agronomic peculiarities of clover to which P.A. Lisitsyn, I.S. Travin, V.D. Shcherbachev, P.A. Sergrrv, the famous Russian scientists, greatly contributed, were initiated in 1930s. Thereat an extensive breeding based on mass selection was mainly used. Since the middle 1950s the research group of clover breeding, and then the Department, this group was transformed into in 1972, were headed by A.S. Novoselova.

Due to theoretical and methodological advance, 25 climatically and ecologically differentiated clover varieties such as VIK 7, Tetraploidnii VIK, Mars, Rannii 2, Trio, Altyn, Topas, Dobrynya, VIK 77, TOS 870, Orlik, VIK 84, Stodolich, Ratibor, Mariya, Pamyati Lisitsyna, Pamyati Burlaca (*Trifolium pratense*); Yubileinii, VIK 70, Lugovik (*Trifolium repens*); Pervenets, Mayak (*Trifolium hybridum*) were derived in this Department, recently headed by M.Yu. Novoselov. Chemical mutagenesis and experimental polyploidy were basically used. These varieties are effective in the northern Russian regions providing dry weight mass yield of 10-12 t per 1 ha. Ultra early ripening clover varieties contribute to expansion of sustainable seed production zone 300-350 km to the north and up to 70 km to the east with the seed yields of 306 centner per 1 ha, being enough to meet the demand on seeds in Russia and to sell abroad to Belarus and Germany (1-4). The Topaz variety obtained by edaphic selection is tolerant to soil acidification (pH 4.5-5.5) and produces the dry weight biomass and seeds at 10-11 t per 1 ha and 2-3 centner per 1 ha, respectively (1).

Our priority lies in scientific, methodological and management interrelations of the experts from 14 institutions taking into account varying soil and climatic conditions on the Russia territory. The monograph «Ecological breeding of clover» (Moscow, 2012) summarizes the results of these complex investigations. In 1999 the creation of the varieties of red clover basic for sustainable forage production and agriculture biologization in Chernozem zone of Russia was awarded the State Prize of the Russian Federation in Science and Technology.

Alfalfa is an important leguminous fodder plant in the world. Nevertheless, it was atypical in the Russian Nechernozem'e under relatively low temperatures, specific water regime and the acidic podzol soils. Due to theoretical experiments initiated in 1930s, in 1956 the Severnaya Gibrinaya variety capable to accumulate enough biomass but, unfortunately, failed to produce mature seeds has been originated by A.M. Konstantinova. Further a new generation of the alfalfa varieties was obtained, namely Vega 87, Lada, Pastbishnaya 88, Lugovaya 67, Selena, Soleustoichivaya, Sonata, Nadezhda, Nakhodka, Galiya, Agniya, which are specific with respect to their environmental, phytocenotic, edaphic traits and agrotechnologies. Vega variety with seed productivity at 2-4 centner per 1 ha was the first variety capable to reproduce successfully in the Central Nechenozem zone. Using phytocenotic selection the Pastbishnaya 88, Lugovaya 62, Nakhodka varieties were bred which in combination with cereal grasses guarantee the dry weight yield of fodder mass at 11-13 t/ha and up to 2.5 t/ha protein output. A phytocenotic longevity is the main peculiarity of these varieties, as after 4-5 year exploitation the ratio of leguminous component is not less than 35-45 % (5, 6). According to L.G. Ramenskii (9), there is a threshold for prosperity of the species when its number is enough competitive. With respect to the competitiveness the Pastbishnaya and Lugovaya varieties overcome the threshold for prosperity being sustainable botanical species in polycomponent agrophytocenosis. Alfalfa edaphic breeding also has been developed, and the obtained Selena plants can produce 10-12 t/ha dry mass on acidic soils at pH 4.5-5.6.

Symbiotic technologies are the promising strategy as it allows improving the nitric and phosphorus plant nutrition due to the plant-microbe interaction.

This approach based on integration of the genetic systems of plants and microorganisms is successfully being developed (7, 8), and the methods for isolation, selection and estimation of rhizobia strains have been patented. Symbiotic selection was used to design the integrated variety-and-microorganism effective systems such as the alfalfa Vega, Lugovaya 67, Pstbishchnaya varieties and 415b rhizobia strain, and Selena and Agniya varieties with 404b strain, and the clover varieties with local K-18 strain. These symbiotic systems allow to harvest 12-14 t dry matter and 2.0-2.5 t protein per 1 ha and accumulate biological nitrogen in soil of 150-200 kg/ha due to roots and crop residues. In the terms of ecology such systems are a biogeocenotic formation at a supraorganismal level denoted by botanist L.G. Rameskii in 1925 (9) and zoologist V.N. Beklemishev in 1951 (10) as consortia which are the primary units of self-organized and self-sustainable agroecosystems sufficient in nitrogen and partly in phosphorus nutrition.

Due to the developed theoretical bases and methodology the varieties of annual legumes with different climatic and environmental features and economic specialization have been originated. They are the spring vetch Lugovskaya 83, Lugovskaya 85, Lugovskaya 98, Vera, Lugovchaanka, Lugovskaya 24, Vakentina, Neposeda and the winter vetch Lugovskaya 2 (11). Many of them are early ripening and produce seeds at 3.0-4.0 t/ha over 65-88 day vegetation. Besides, they are less heat-loving, thus being successfully cultivated in regions with insufficient heat and excessive soil moisture. Yu.S. Tyuring has suggested creating a specialized forage vetch, and the Lugovskaya 98 variety is now widely used.

Perennial forage grasses and landscape are major components of the hay and pasture agro-ecosystems, so more than 20 varieties of perennial grasses were zoned. They are smooth brome-grass Fakelnii and Morshanets, timothy grass VIK 85, meadow fescue Krasnopoimskaya 92, tall fescue Lira, cocksfoot Morshanskaya 89, perennial ryegrass Tetraploidnii, VIK 66 and Duet, meadow grass Pobeda, Tambovets, fescue-ryegrass hybrid VIK 90 (*Festulolium*). They produce 11-12 t/ha dry matter, are more nutritionally valuable and more tolerant to diseases (12, 13).

Drought and salt tolerant fodder xerohalophytes are used on the arid Russia territories, i.e. kochia Barkhan, Dzhangar, Delta and Istok varieties, eurotia Favorite and Bar varieties, salsola Salang variety, *Camphorosma lessingii* Alsu and Nogana varieties, buckwheat Tsag variety, artemisia Sonet variety, and licorice Fortuna variety, which were obtained in cooperation with the researchers from A.N. Kostyakov Institute of Hydro Engineering and Land Reclamation (Moscow) and Kalmykia Research Institute of Agriculture (Elista). These fodder crops are widely used in degraded dry steppes and semi-steppes to increase productivity of the pastures (14-16).

Collecting gene pool together with application of biotechnology and investigations on plant immunity helpful to create tolerant initial material are being the basic approaches (17, 18). In 1930s the mobilization of fodder plant genetic recourses began. In more than 45 expeditions held in cooperation with the N.I. Vavilov Research Institute of Plant Industry (VIR, St. Petersburg), over 6 000 specimens of the wild grass seeds have been sampled in different regions. As a result, valuable alfalfa, clover, sweet clover, sainfoin, rump, foxtail, wheatgrass, wheatgrass forms, as well as endemics have been found, and this work is ongoing. Currently the gene pool collection includes 447 species and about 6500 specimens (16). For clover the polymorphic molecular markers are patented (19). Two clover genetic maps are built in cooperation with Japanese researchers.

Biotechnologies were used to create promising forms for breeding (20-23) such as the clover tolerant to soil acidity, the transgenic clove and alfalfas regenerants, the breeding alfalfa forms tolerant to salination. Using extremely high

tolerance to salinity the alfalfa *Soleustoichivaya* variety was obtained which is successful on the secondary salinized soils in Nizhnee Povolzhie.

In view of disease-resistant fodder plant development the infectious agents were indicated with identification of those the most aggressive. The procedure for cultivation of these pathogens was suggested. Using field test the tolerance was estimated in more than 3500 plants. The disease resistant plants were derived by subjecting to high levels of disease pressure with further recurrent selection, hybridization and polycrossing. As a result the clover, smooth brome-grass and timothy grass resistant to diseases have been developed (24).

Generally, plant tolerance to stresses, the selection of disease-resistant forms, genetic transformation and regeneration, and finding microsymbionts of the selected plants were under consideration (25-38).

Much attention was paid to seed production of fodder crops such as fescue *Krasnopoimskaya* 92 and *Kvarta* varieties, hairy vetch, alfalfa *Sonet* and *Tambovchanka* varieties, red fescue *Diana* and *Sigma* varieties, *festulolum* VIK 90 and *Izumrudnii* varieties, meadow grass *Tambovets* variety.

The location of cultivation areas are revised for clover seed reproduction, for cultivation and commercial reproduction of alfalfa perennial ryegrass, timothy, meadow fescue, cocksfoot and smooth brome-grass, and 16 parameters are currently developed for seed quality assessment.

The leading position of the Center is due to wide research cooperation with other Russian researchers and the foreign scientists from Kazakhstan, Uzbekistan, USA, Israel, Belarus, and Mongolia. Thus, the mobilization of genetic resources and breeding fodder crops is particularly carried out together with *Norddeutsche Pflanzenzucht Hans-Georg Lembke KG* (Germany). The breeders of the Fodder Research Institute were recognized the leaders in Russia in developing botanic, geographical, environmental and evolutionary bases for breeding fodder plants to manage a sustainable forage production.

The fundamental aspects of modern strategy for current fodder crop breeding and seed reproduction are based on mobilizing genetic resources of both cultivated and wild plants, using data banking systems, collecting genetic resource as the donors of desired traits, QTLs marking and genetic mapping in clover and other valuable fodder plants, designing gene constructions to develop transgenic plants resistant to fungal pathogens and environmental stresses. An induced re-combinogenesis and mutagenesis, as well as interspecial and intraspecial interrelations of the plants different in adaptive strategy are also helpful to design the parental forms by means of the improved phytocenotic selection. One more edaphic approach involves plant adaptation to acidic and salinized soils and toxic chlorine, manganese and sodium ions. As far as the plant-microbe interactions are related to fodder plant productivity, it necessitates the study of their efficacy during plant ontogenesis as influenced by the genetic traits and geographic origin of the partners. To substantiate the principles and approaches to preemptive breeding disease-resistant plants, it is essential to forecast changes of a microbial landscape. The study of reproductive peculiarities of developed or introduced fodder plants is helpful with the view of optimization of seed reproduction. Both global and regional survey of climatic changed and their impact allows to suggest the strategy for breeding climatically and edaphically specialized fodder crops.

Applied researches are focused on minimization of the global deficit in resources and the negative effects of modern agriculture. Thus, development of the complementary varieties with climatic and environmental specialization provides for a sustainable livestock forage production subject to the principles of ecological agriculture. Thus the clover, alfalfa and grassland legumes with different economic specialization, high green mass and seed yield, enhanced nitrogen

fixation and an increased resistance to most diseases and abiotic factors should be developed. Using biogeocenotic and improved breeding techniques the perennial cereal grasses with high forage value, disease-resistant, tolerant to abiotic stresses should be obtained to be used for mowing and grazing. Phytocenotic selection and induced recombinogenesis are applied to create spring and winter vetch for different economic purposes. An adaptive selection is suggested for cole fodder crops such as rape, colza, etc., with high yield, increased nutritive value and tolerant to biotic agents and edaphic factors. Arid fodder plants with specific edaphic and phytocenotic traits and more drought- and salt-tolerant should be developed for restoration of zonal biodiversity and fertility in soils after salination and in the degraded pastures. Environmental and phytocenotic approach is basic in agro ecological seed reproduction and more effective realization and maintenance of plant genetic potential. It is also helpful in improving specific agro technology for each variety in view of anticipated climatic changes and also at estimation of germination of seeds in fodder plant species and varieties.

Environmental and evolutionary principles are basic in both fundamental and applied studies. This approach at studying biological phenomena, particularly appearance and development of adaptive capability during phylo- and ontogenesis has been suggested by A.A. Zhuchenko in the monographs «Ecological genetics of cultivated plants: adaptation, recombinogenesis, agrobiocenosis» (Moscow, 1980) and «Ecological genetics of cultivated plants as a scientific discipline» (Moscow, 2010).

We suggested a breeding paradigm based on fundamental biogeocenotic principles of fodder crop breeding and seed production as a part of adaptive plant cultivation (39-41). According to the paradigm, the phototrophic fodder plants trigger the flow of organic matter and energy in soil, resulting in formation of a biotic complex, the fodder agro biocenosis, being a system of functionally related components. In the system a phytocenotic interaction between the plants, an edaphic interaction between plants and soil, and plant-microbe symbiotic and/or associative interaction are dominant. These interactions are the breeding potential resources which are essential for design the fodder crops as a biocenotic units possessing phytocenotic, edaphic, symbiotic and ecotypic specificity enough to form an adaptive self-organizing sustainable and high-yielding structures. A capability to self-optimization in these systems is functionally irreplaceable and economically valuable. Therefore, the biogeocenotic breeding of fodder crops can be considered as special scientific discipline.

Thus, more than 150 fodder crop varieties developed for past 42 years and widely cultivated in Russia are not inferior to the foreign varieties on productivity but more frost resistant, edaphic tolerant to soil acidity and salination and compatible in mixtures. More than 40 varieties of perennial grasses, specifically smooth brome-grass, timothy grass, meadow fescue, cocksfoot, perennial ryegrass, meadow grass, with increased productivity above 11-12 tons per hectare, higher disease resistance and improved nutritive value are developed. For restoration of biodiversity and soils in degraded pastures of the arid southern regions of Russia 18 varieties of fodder xenohalophytes are suggested such as *Kochia prostrata* (L.) Schrad. Barkhan and Dzhabgar varieties, *Salsola orientalis* S.G. Gmel. Salang variety, *Camphorosma lessingii* (Litv.) Nogana and Alsuv varieties, *Eurotia ceratoides* (L.) C.A. Mey. Favorit, Tulkin and Bar varieties, etc., widely used for ecological restoration of desertified soils. The applied researches are influenced by recent state of agriculture, global deficit of natural resources and climate changes and based on fundamental research in genetics, gene engineering, biotechnology, botany, ecology, etc., elucidating expression of valuable traits and their inheritance in a sustainable developing population.

As a conclusion the paradigm is suggested based on the fundamental biogeocenotic approaches and principles of adaptive plant cultivation, particularly breeding and seed production technologies. The biogeocenotic breeding of fodder crops can be considered as special scientific discipline.

REFERENCES

1. Novoselov M.Yu. *Seleksiya klevra lugovogo (Trifolium pratense L.)* [Breeding of *Trifolium pratense* L.]. Moscow, 1999.
2. Novoselov M.Yu. V sbornike: *Adaptivnaya sistema seleksii kormovykh rastenii (biogeotsenoticheskii podkhod)* [In: Forage plants: adaptive breeding and biogeocenotic approach]. Moscow, 2007: 65-69.
3. Novoselov M.Yu. V sbornike: *Ekologicheskaya seleksiya i semenovodstvo klevra lugovogo* [In: Ecological selection and seed reproduction of *Trifolium pratense* L.]. Moscow, 2012: 22-55.
4. Novoselova A.S., Novoselov M.Yu. V sbornike: *Ekologicheskaya seleksiya i semenovodstvo klevra lugovogo* [In: Ecological selection and seed reproduction of *Trifolium pratense* L.]. Moscow, 2012: 7-21.
5. Piskovatskii Yu.M., Nenarokov Yu.M., Solozhentseva L.F. V sbornike: *Adaptivnaya sistema seleksii kormovykh rastenii (biogeotsenoticheskii podkhod)* [In: Forage plants: adaptive breeding and biogeocenotic approach]. Moscow, 2007: 30-63.
6. Piskovatskii Yu.M., Nenarokov Yu.M., Stepanova G.V. V sb.: *Adaptivnoe kormoproizvodstvo: problemy i resheniya* [In: Adaptive fodder production: problems and solutions]. Moscow, 2002: 294-308.
7. Drobysheva L.V., Zyatchina G.P. V sbornike: *Kormoproizvodstvo: problemy i puti resheniya* [In: Fodder production: effective technologies]. Moscow, 2007: 262-270.
8. Stepanova G.V., Nizhnik O.V., Selitskaya O.V., Antonova L.S. V sbornike: *Kormoproizvodstvo: problemy i puti resheniya* [In: Fodder production: effective technologies]. Moscow, 2007: 357-364.
9. Ramenskii L.G. *Vvedenie v kompleksnoe pochvenno-geobotanicheskoe issledovanie zemel'* [Introduction to a comprehensive soil and geobotanical study land]. Moscow, 1925.
10. Beklemishev V.N. *Byulleten' MOIP, Otdel Biol.*, 1951, 56(6): 53-65.
11. Tyurin Yu.S. V sbornike: *Kormoproizvodstvo: problemy i puti resheniya* [In: Fodder production: effective technologies]. Moscow, 2007: 306-310.
12. Kuleshov G.F., Bekhtin N.S., Klochkova V.S., Malyuzhenets E.E., Morozova E.A. V sbornike: *Kormoproizvodstvo: problemy i puti resheniya* [In: Fodder production: effective technologies]. Moscow, 2007: 294-301.
13. Kostenko S.I., Kuleshov G.F., Bekhtin N.S., Klochkova V.S., Morozova E.A. V sbornike: *Kormoproizvodstvo: problemy i puti resheniya* [In: Fodder production: effective technologies]. Moscow, 2007: 301-306.
14. Balnokin Yu.V., Myasoedov N.A., Shamsutdinov Z.Sh., Shamsutdinov N.Z. *Fiziologiya rastenii*, 2007, 52(6): 282-290.
15. Shamsutdinov Z.Sh. *Kormoproizvodstvo*, 2010, 8: 25-27.
16. Shamsutdinov Z.Sh., Savchenko I.V., Shamsutdinov N.Z. *Vestnik Rossiiskoi akademii sel'skokhozyaistvennykh nauk*, 2006, 3: 79-81.
17. Kozlov N.N., Korovina V.L., Makarenkov M.A., Klimenko I.A., Trukhan V.A., Komkova T.N. V sbornike: *Kormoproizvodstvo: problemy i puti resheniya* [In: Fodder production: effective technologies]. Moscow, 2007: 335-343.
18. Shamsutdinov Z.Sh. *Doklady Rossiiskoi akademii sel'skokhozyaistvennykh nauk*, 2002, 3: 28-30.
19. Kozlov N.N., Pribytkova T.F., Malysheva Yu.N. *Sposob markirovaniya selektsionnykh dostizhenii klevra lugovogo na osnove RAPD-markerov. Patent № 2244416 (RF). Opubl. 20.01.2005. Byul. № 2* [Use of RAPD markers to identify the products of *Trifolium pratense* L. breeding achievements. Patent RU № 2244416. Published 01.20.2005. Bulletin № 2].
20. Klochkova V.S., Korovina V.L., Kuleshov G.F. *Timofeevka lugovaya VIK 85. Patent na selektsionnoe dostizhenie № 0440. Opubl. 29.11.1999* [Timothy grass VIK 85. Patent for breeding achievements (Russia) № 0440. Published 11.29.1999].
21. Ostretsova I.N., Mazur K.K., Novoselov M.Yu., Mazin V.V., Solozhentsev P.D. *Sposob polucheniya triploidnykh rastenii lyutserny. Patent № 1595408 (SSSR). Opubl. 04.07.1988. Byul. № 36* [Technology used to obtain triploids of alfalfa plants. Patent SU № 1595408. Published 07.04.1988. Bulletin № 36].
22. Solozhentsev P.D., Solozhentseva L.F., Agafodorova M.N. *Sposob povysheniya ustoichivosti rastenii lyutserny k fuzariozu. Patent № 2278508 (RF). Opubl. 27.06.2006. Byul. № 18* [Technology used to increase tolerance to fusariosis in alfalfa plants. Patent RU № 2278508. Published 06.27.2006. Bulletin № 18].
23. Mezentsev A.V., Lyubavina L.A. *Sposob regeneratsii rastenii klevra. A.S. № 888861 (SSSR). Opubl. 15.12.1981. Byul. № 46* [Method for clover plant regeneration. A.S. SU № 888861. Published 12.15.1981. Bulletin № 46].

24. Putsa N.M., Razgulyaeva N.V., Kostenko N.Yu. V sbornike: *Kormoproizvodstvo: problemy i puti resheniya* [In: Fodder production: effective technologies]. Moscow, 2007: 371-377.
25. Ivashuta S.I., Agafodorova M.N., Mazin V.V. *Sposob regeneratsii rastenii lyutserny in vitro*. Patent № 2073426 (RF). Opubl. 20.02.1997. Byul. № 5 [Method for alfalfa plant regeneration in vitro. Patent SU № 2073426. Published 02.20.1997. Bulletin № 5].
26. Stepanova G.V., Piskovatskii Yu.M., Nenarokov Yu.M. *Sposob imitatsii vytyapyvaniya mnogoletnikh trav zivotnymi*. A.S. № 1709965 (SSSR). Opubl. 07.02.1992. Byul. № 5 [Method to simulate perennial grass trampling by animals. A.S. SU № 1709965. Published 02.07.1992. Bulletin № 5].
27. Zolotarev V.N. *Sposob proizvodstva semyan ul'trarannespelogo klevera lugovogo i timofeevki lugovoi*. Patent № 2216918 (RU). Opubl. 27.11.2003. Byul. № 33 [Method for seed production in ultra early ripening clover and timothy grass. Patent RU № 2216918. Published 11.27.2003. Bulletin № 33].
28. Mazur K.K., Smurygin M.A., Ostretsova I.N., Zarutskaya L.S., Novoselov M.Yu., Oshanina N.P. *Sposob vyrashchivaniya klevera lugovogo v zashchishchennom grunte*. A.S. № 1184474 A (SSSR). Opubl. 15.10.1985. Byul. № 38 [Technology of clover growing in green houses. A.S. SU № 1184474 A. Published 10.15.1985. Bulletin № 38].
29. Stepanova G.V. *Sposob seleksii lyutserny*. Patent № 2077190 (RF). Opubl. 20.04.1997. Byul. № 11 [Method of alfalfa breeding. Patent RU № 2077190. Published 04.20.1997. Bulletin № 11].
30. Zolotarev V.N. *Sposob polucheniya semyan diploidnogo klevera lugovogo ul'tra rannespelogo tipa*. Patent № 2226048 (RU). Opubl. 27.03.2004. Byul. № 9 [Method of seed production in diploid plants of ultra early ripening clover. Patent RU № 2226048. Published 03.27.2004. Bulletin № 9].
31. Novoselov M.Yu., Paivin S.G. *Sposob otbora morozostoikikh form klevera lugovogo*. Patent № 2091011 (RF). Opubl. 27.09.1997. Byul. № 27 [Method of selection of frost resistant clover plants. Patent RU № 2091011. Published 09.27.1997. Bulletin № 27].
32. Novoselov M.Yu., Paivin S.G. *Sposob otbora kisloutostoichivyykh form klevera lugovogo*. Patent № 2138154 (RF). Opubl. 27.09.1999. Byul. № 10 [Method of selection of clover plants resistant to acidic conditions. Patent RU № 2138154. Published 09.27.1999. Bulletin № 10].
33. Zolotarev V.N., Seregin S.V. *Sposob podbora komplementarnykh kul'tur dlya smeshannykh posevov*. Patent № 2306687 (RU). Opubl. 27.09.2007. Byul. № 27 [Method of selection of complementary crops to compose their mixtures for cultivation. Patent RU № 2306687. Published 09.27.2007. Bulletin № 27].
34. Zolotarev V.N., Krasavina N.Yu. *Sposob podbora komplementarnogo po allelopaticeskomu posledeystviyu predshestvennika*. Patent № 2436285 (RU). Opubl. 20.12.2011. Byul. № 35 [Method of selection of a precursor with complementary allelopathic aftereffect. Patent RU № 2436285. Published 12.20.2011. Bulletin № 35].
35. Solozhentsev P.D. *Sposob vydeleniya komponentov iz pyl'tsy pokrytosemennyykh rastenii*. A.S. № 1145954 A (SSSR). Opubl. 23.03.1985. Byul. № 11 [Method of component extraction from the pollen of angiosperms. A.S. SU № 1145954 A. Published 03.23.1985. Bulletin № 11].
36. Mazur K.K., Smurygin M.A., Ostretsova I.N., Serebryakova L.E., Zarutskaya L.S., Oshanina N.P., Mel'nikov E.V., Novoselov M.Yu., Letyagin Yu.M. *Sposob vyrashchivaniya lyutserny v kontroliruemyykh usloviyakh*. A.S. № 933053 (SSSR). Opubl. 07.06.1982. Byul. № 21 [Technology of alfalfa growing under controlled conditions. A.S. SU № 933053. Published 06.07.1982. Bulletin № 21].
37. Mazur K.K., Smurygin M.A., Zarutskaya L.S., Oshanina N.P., Ostretsova I.N., Novoselov M.Yu., Solozhentsev P.P., Kozik L.A. *Sposob vyrashchivaniya kostretsna bezostogo v zashchishchennom grunte*. A.S. № 1416086 (SSSR). Opubl. 15.08.1988. Byul. № 30 [Technology of smooth brome-grass growing in green houses. A.S. SU № 1416086. Published 08.15.1988. Bulletin № 30].
38. Novoselov M.Yu., Mazur K.K., Oshanina N.P., Ostretsova I.N., Zarutskaya L.S. *Sposob polucheniya mutantov i miksploidov klevera lugovogo v zashchishchennom grunte*. A.S. № 1470251 A1 (SSSR). Opubl. 07.04.1989. Byul. № 13 [Method to obtain mutants and mixoploids of clover in green houses. A.S. SU № 1470251 A1. Published 04.07.1989. Bulletin № 13].
39. Shamsutdinov Z.Sh. *Selektsiya i semenovodstvo*, 2004, 3: 2-10.
40. Shamsutdinov Z.Sh. *Kormoproizvodstvo*, 2007, 5: 24-27.
41. Shamsutdinov Z.Sh., Piskovatskii Yu.M., Novoselov M.Yu. *Kormoproizvodstvo*, 2008, 1: 2-6.